

Zeta-functions for germs of meromorphic functions and Newton diagrams

S.M. Gusein-Zade I. Luengo A. Melle-Hernández *

Abstract

For a germ of a meromorphic function $f = \frac{P}{Q}$, we offer notions of the monodromy operators at zero and at infinity. If the holomorphic functions P and Q are non-degenerated with respect to their Newton diagrams, we give an analogue of the formula of Varchenko for the zeta-functions of these monodromy operators.

1 Germs of meromorphic functions

A polynomial f of $(n+1)$ complex variables of degree d determines a meromorphic function f on $\mathbb{CP}^{n+1}$. If one wants to understand the behaviour of f at infinity, it is natural to analyze germs of the meromorphic function f at points from the infinite hyperplane $\mathbb{CP}_\infty^n \subset \mathbb{CP}^{n+1}$. In local analytic coordinates $z_0, z_1, \dots, z_n$, centred at a point $p \in \mathbb{CP}_\infty^n$ such that the infinite hyperplane $\mathbb{CP}_\infty^n$ is given by the equation $\{z_0 = 0\}$, the germ of the function f has the form $f = \frac{P(z_0, \dots, z_n)}{z_0^d}$. Let us consider germs of meromorphic functions of a general form.

DEFINITION 1 A *germ of a meromorphic function on $(\mathbb{C}^{n+1}, 0)$* is a fraction $f = \frac{P}{Q}$, where P and Q are germs of holomorphic functions $(\mathbb{C}^{n+1}, 0) \rightarrow (\mathbb{C}, 0)$. Two germs of meromorphic functions $f = \frac{P}{Q}$ and $f' = \frac{P'}{Q'}$ are *equal* if there exists a germ of a holomorphic function $U : (\mathbb{C}^{n+1}, 0) \rightarrow \mathbb{C}$ such that $U(0) \neq 0$, $P' = U \cdot P$ and $Q' = U \cdot Q$.

REMARKS. (1) For our convenience here we do not consider functions of the type $\frac{1}{Q(z)}$ or $\frac{P(z)}{1}$.

(2) According to the definition $\frac{x}{y} \neq \frac{x^2}{xy}$, but $\frac{x}{y} = \frac{x \exp(x)}{y \exp(x)}$.

Recently V.I. Arnold had obtained classifications of simple germs of meromorphic functions for certain equivalence relations.

In what follows we shall consistently use resolutions of germs of meromorphic functions.

*First author is partially supported by RFBR-95-01-01122a and INTAS-4373. Last two authors are partially supported by CAICYT PB94-291.

DEFINITION 2 A *resolution of the germ f* is a modification of $(\mathbb{C}^{n+1}, 0)$ (i.e. a proper analytic map $\pi : \mathcal{X} \rightarrow \mathcal{U}$ of a smooth analytic manifold $\mathcal{X}$ onto a neighbourhood $\mathcal{U}$ of the origin in $\mathbb{C}^{n+1}$, which is an isomorphism outside of a proper analytic subspace in $\mathcal{U}$) such that the total transform $\pi^{-1}(H)$ of the hypersurface $H = \{P = 0\} \cup \{Q = 0\}$ is a normal crossing divisor at each point of $\mathcal{X}$.

The fact that the preimage $\pi^{-1}(H)$ is a divisor with normal crossings implies that in a neighbourhood of any point of it, there exists a local system of coordinates $y_0, y_1, \dots, y_n$ such that the liftings $\tilde{P} = P \circ \pi$ and $\tilde{Q} = Q \circ \pi$ of the functions P and Q to the space $\mathcal{X}$ of the modification are equal to $u y_0^{k_0} y_1^{k_1} \dots y_n^{k_n}$ and $v y_0^{l_0} y_1^{l_1} \dots y_n^{l_n}$ respectively, where $u(0) \neq 0$ and $v(0) \neq 0$.

Let B_ε be the closed ball of radius ε with the centre at the origin in $\mathbb{C}^{n+1}$ and ε be small enough such that (representatives of) the functions P and Q are defined in B_ε and for any positive $\varepsilon' < \varepsilon$ the sphere $S_{\varepsilon'} = \partial B_{\varepsilon'}$ intersects the analytic spaces $\{P = 0\}$, $\{Q = 0\}$ and $\{P = Q = 0\}$ transversally (in the stratified sense). We choose δ small enough and take the ball $B_\delta \subset \mathbb{C}^2$ of radius δ centred at the origin.

DEFINITION 3 Let $c \in \mathbb{C}$ be such that $\|c\|$ is small enough, the *0-Milnor fibre $\mathcal{M}_f^0$ of the germ f* is the set

$$\mathcal{M}_f^0 = \{z \in B_\varepsilon : (P(z), Q(z)) \in B_\delta \subset \mathbb{C}^2, f(z) = \frac{P(z)}{Q(z)} = c\}.$$

In the same way, for $c \in \mathbb{C}$ such that $\|c\|$ is large enough, the *∞ -Milnor fibre $\mathcal{M}_f^\infty$ of the germ f* is the set

$$\mathcal{M}_f^\infty = \{z \in B_\varepsilon : (P(z), Q(z)) \in B_\delta \subset \mathbb{C}^2, f(z) = \frac{P(z)}{Q(z)} = c\}.$$

LEMMA 1 *The notion of the 0- (respectively of the ∞ -) Milnor fibre is well defined, i.e. for $\|c\|$ small enough: $\|c\| \ll \delta \ll \varepsilon$ (respectively for $\|c\|$ large enough: $\|c\|^{-1} \ll \delta \ll \varepsilon$) the differentiable type of $\mathcal{M}_f^0$ (respectively of $\mathcal{M}_f^\infty$) does not depend on ε, δ and c .*

Proof. Let $\pi : \mathcal{X} \rightarrow \mathcal{U}$ be a resolution of the germ f which is an isomorphism outside the hypersurface $H = \{P = 0\} \cup \{Q = 0\}$. Let $r : \mathbb{C}^{n+1} \rightarrow \mathbb{R}$ be the function $r(z) = \|z\|^2$, let $\tilde{r} = r \circ \pi : \mathcal{X} \rightarrow \mathbb{R}$ be the lifting of r to the space $\mathcal{X}$ of the resolution. For ε small enough, the hypersurface $\tilde{S}_\varepsilon = \{\tilde{r} = \varepsilon^2\}$ (the preimage of the sphere $S_\varepsilon \subset \mathbb{C}^{n+1}$) is transversal to all components of the total transform $\pi^{-1}(H)$. At each point of $\pi^{-1}(H)$ in a local coordinate system one has $P \circ \pi = u y_0^{k_0} \dots y_n^{k_n}$, $Q \circ \pi = v y_0^{l_0} \dots y_n^{l_n}$ with $u(0) \neq 0$ and $v(0) \neq 0$. Thus $f \circ \pi = w y_0^{m_0} \dots y_n^{m_n}$ with $w(0) \neq 0$. The real hypersurface $\tilde{S}_\varepsilon$ is transversal to all coordinate subspaces (of different dimensions). It is not difficult to show that this implies transversality of $\tilde{S}_\varepsilon$ to the (complex) hypersurfaces $\{w y_0^{m_0} \dots y_n^{m_n} = c\}$ for $\|c\|$ small enough and for $\|c\|$ large enough. Now the proof follows from the standard arguments.

REMARKS. (1) The definition means that $\mathcal{M}_f^0$ or $\mathcal{M}_f^\infty$ is equal to

$$\{z \in B_\varepsilon : (P(z), Q(z)) \in B_\delta \subset \mathbb{C}^2, P(z) = cQ(z), \quad P(z) \neq 0\}$$

and thus the Milnor fibres of the functions $\frac{P}{Q}$ and $\frac{RP}{RQ}$ with $R(0) = 0$ are, generally speaking, different.

(2) For $f = \frac{P}{Q}$, let $f^{-1} = \frac{Q}{P}$. It is not difficult to understand that $\mathcal{M}_{f^{-1}}^0 = \mathcal{M}_f^\infty$ and $\mathcal{M}_{f^{-1}}^\infty = \mathcal{M}_f^0$. Just the same properties hold for the monodromy transformations and for the zeta-functions discussed below.

(3) It is possible (and sometimes more convenient) to define the Milnor fibres as follows:

$$\mathcal{M}_f^0 = \{z \in B_\varepsilon : \|Q(z)\| \leq \delta, P(z) = cQ(z) \neq 0\}$$

with $\|c\| \ll \delta \ll \varepsilon$, and

$$\mathcal{M}_f^\infty = \{z \in B_\varepsilon : \|P(z)\| \leq \delta, P(z) = cQ(z) \neq 0\}$$

with $\|c\|^{-1} \ll \delta \ll \varepsilon$.

The meromorphic function f determines a map from $B_\varepsilon \setminus \{P = Q = 0\}$ to the projective line $\mathbb{CP}^1$ ($z \mapsto (P(z) : Q(z))$), which also will be denoted by f . Lemma 1 implies that this map is a locally trivial fibration in punctured neighbourhoods of the points $0 = (0 : 1)$ and $\infty = (1 : 0)$ of $\mathbb{CP}^1$.

DEFINITION 4 The *0-monodromy transformation* h_f^0 (respectively the *∞ -monodromy transformation* h_f^∞) of the germ f is the monodromy transformation of the fibration f over the loop $c \cdot \exp(2\pi it)$, $t \in [0, 1]$, with $\|c\|$ small enough (respectively large enough).

The 0- or ∞ - monodromy operator is the action of the corresponding monodromy transformation in a homology group of the Milnor fibre. We are interested to apply the results for meromorphic functions to the problem of calculating the zeta-function of a polynomial at infinity. Thus we shall consider the zeta-functions $\zeta_f^0(t)$ and $\zeta_f^\infty(t)$ of the corresponding monodromy transformations:

$$\zeta_f^\bullet = \prod_{q \geq 0} \{\det[id - t h_{f*}^\bullet|_{H_q(\mathcal{M}_f^\bullet; \mathbb{C})}]\}^{(-1)^q}$$

($\bullet = 0$ or ∞). This definition coincides with that in [2] and differs by minus sign in the exponent from that in [1].

2 Resolution of singularities and the formula of A'Campo for germs of meromorphic functions

Let $f = \frac{P}{Q}$ be a germ of a meromorphic function on $(\mathbb{C}^{n+1}, 0)$ and let $\pi : \mathcal{X} \rightarrow \mathcal{U}$ be a resolution of the germ f . The preimage $\mathcal{D} = \pi^{-1}(0)$ of the origin of $\mathbb{C}^{n+1}$, is a

normal crossing divisor. Let $S_{k,l}$ be the set of points of $\mathcal{D}$ in a neighbourhood of which the functions $P \circ \pi$ and $Q \circ \pi$ in some local coordinates have the form $u y_0^k$ and $v y_0^l$ respectively ($u(0) \neq 0, v(0) \neq 0$). A slight modification of the arguments of A'Campo ([1]) permits to obtain the following version of his formula for the zeta-function of the monodromy of a meromorphic function.

THEOREM 1 *Let the resolution $\pi : \mathcal{X} \rightarrow \mathcal{U}$ be an isomorphism outside the hypersurface $H = \{P = 0\} \cup \{Q = 0\}$. Then*

$$\begin{aligned}\zeta_f^0(t) &= \prod_{k>l} (1 - t^{k-l})^{\chi(S_{k,l})}, \\ \zeta_f^\infty(t) &= \prod_{k<l} (1 - t^{l-k})^{\chi(S_{k,l})}.\end{aligned}$$

REMARK. A resolution π of the germ $f' = \frac{RP}{RQ}$ is at the same time a resolution of the germ $f = \frac{P}{Q}$. Moreover the multiplicities of any component C of the exceptional divisor in the zero divisors of the liftings $(RP) \circ \pi$ and $(RQ) \circ \pi$ of the germs RP and RQ are obtained from those of the germs P and Q by adding one and the same integer, the multiplicity $m = m(C)$ of R . Nevertheless the meromorphic functions f and f' can have different zeta-functions. The reason why formulae in the previous theorem give different results for f and f' consists in the fact that if an open part of the component C lies in $S_{k,l}(f)$ then, generally speaking, its part which lies in $S_{k+m,l+m}(f')$ is smaller.

3 Zeta-functions of meromorphic functions via partial resolutions

Let $f = \frac{P}{Q}$ be a germ of a meromorphic function on $(\mathbb{C}^{n+1}, 0)$ and let $\pi : (\mathcal{X}, \mathcal{D}) \rightarrow (\mathbb{C}^{n+1}, 0)$ be an arbitrary modification of $(\mathbb{C}^{n+1}, 0)$, which is an isomorphism outside the hypersurface $H = \{P = 0\} \cup \{Q = 0\}$ (i.e. π is not necessarily a resolution). Let $\varphi = f \circ \pi$ be the lifting of f to the space of the modification, i.e. the meromorphic function $\frac{P \circ \pi}{Q \circ \pi}$. For a point $x \in \pi^{-1}(H)$, let $\zeta_{\varphi,x}^0(t)$ and $\zeta_{\varphi,x}^\infty(t)$ be the zeta-functions of the 0- and ∞ -monodromies of the germ of the function φ at x . Let $\mathcal{S} = \{\Xi\}$ be a prestratification of $\mathcal{D} = \pi^{-1}(0)$ (that is a partitioning into semi-analytic subspaces without any regularity conditions) such that, for each stratum Ξ of $\mathcal{S}$, the zeta-functions $\zeta_{\varphi,x}^0(t)$ and $\zeta_{\varphi,x}^\infty(t)$ do not depend on x , for $x \in \Xi$. We denote this zeta-functions by ζ_Ξ^0 and by ζ_Ξ^∞ respectively. The same arguments which were used in [4] imply

THEOREM 2 *For $\bullet = 0$ or ∞ ,*

$$\zeta_f^\bullet(t) = \prod_{\Xi \in \mathcal{S}} [\zeta_\Xi^\bullet(t)]^{\chi(\Xi)}.$$

4 Zeta-functions via Newton diagrams

For a germ $R = \sum a_k x^k : (\mathbb{C}^{n+1}, 0) \rightarrow (\mathbb{C}, 0)$ of a holomorphic function ($k = (k_0, k_1, \dots, k_n)$, $x^k = x_0^{k_0} x_1^{k_1} \dots x_n^{k_n}$), its Newton diagram $\Gamma = \Gamma(R)$ is the union of the compact faces of the polytope $\Gamma_+ = \Gamma_+(R) = \text{convex hull of } \bigcup_{k: a_k \neq 0} (k + \mathbb{R}_+^{n+1}) \subset \mathbb{R}_+^{n+1}$.

Let $f = \frac{P}{Q}$ be a germ of a meromorphic function on $(\mathbb{C}^{n+1}, 0)$ and let $\Gamma_1 = \Gamma(P)$ and $\Gamma_2 = \Gamma(Q)$ be the Newton diagrams of P and Q . We call the pair $\Lambda = (\Gamma_1, \Gamma_2)$ of Newton diagrams Γ_1 and Γ_2 the *Newton pair of f* . We say that the germ of the meromorphic function f is *non-degenerated with respect to its Newton pair $\Lambda = (\Gamma_1, \Gamma_2)$* if the pair of germs (P, Q) is non-degenerated with respect to the pair $\Lambda = (\Gamma_1, \Gamma_2)$ in the sense of [7] (which is an adaptation for *germs* of complete intersections of the definition of A.G. Khovanskii, [5]).

Let us define zeta-functions $\zeta_\Lambda^0(t)$ and $\zeta_\Lambda^\infty(t)$ for a Newton pair $\Lambda = (\Gamma_1, \Gamma_2)$. Let $1 \leq l \leq n+1$ and let $\mathcal{I}$ be a subset of $\{0, 1, \dots, n\}$ with the number of elements $\#\mathcal{I}$ equal to l . Let $L_{\mathcal{I}}$ be the coordinate subspace $L_{\mathcal{I}} = \{k \in \mathbb{R}^{n+1} : k_i = 0 \text{ for } i \notin \mathcal{I}\}$ and $\Gamma_{i, \mathcal{I}} = \Gamma_i \cap L_{\mathcal{I}} \subset L_{\mathcal{I}}$. Let $L_{\mathcal{I}}^*$ be the dual of $L_{\mathcal{I}}$ and $L_{\mathcal{I}+}^*$ the positive orthant of it (the set of covectors which have positive values on $L_{\mathcal{I} \geq 0} = \{k \in L_{\mathcal{I}} : k_i \geq 0 \text{ for } i \in \mathcal{I}\}$). For a primitive integer covector $a \in (\mathbb{R}^*)_+^{n+1}$, let $m(a, \Gamma) = \min_{x \in \Gamma} (a, x)$ and let $\Delta(a, \Gamma) = \{x \in \Gamma : (a, x) = m(a, \Gamma)\}$. We denote by $m_{\mathcal{I}}$ and $\Delta_{\mathcal{I}}$ the corresponding objects for the diagram $\Gamma_{\mathcal{I}}$ and a primitive integer covector $a \in L_{\mathcal{I}+}^*$. Let $E_{\mathcal{I}}$ be the set of primitive integer covectors $a \in L_{\mathcal{I}+}^*$ such that $\dim(\Delta(a, \Gamma_1) + \Delta(a, \Gamma_2)) = l-1$ (the Minkowski sum $\Delta_1 + \Delta_2$ of two polytopes Δ_1 and Δ_2 is the polytope $\{x = x_1 + x_2 : x_1 \in \Delta_1, x_2 \in \Delta_2\}$). There exists only a finite number of such covectors. For $a \in E_{\mathcal{I}}$, let $\Delta_1 = \Delta(a, \Gamma_1)$, $\Delta_2 = \Delta(a, \Gamma_2)$ and

$$V_a = \sum_{s=0}^{l-1} V_{l-1}(\underbrace{\Delta_1, \dots, \Delta_1}_{s \text{ terms}}, \underbrace{\Delta_2, \dots, \Delta_2}_{l-1-s \text{ terms}}),$$

where the definition of the (Minkowski) mixed volume $V(\Delta_1, \dots, \Delta_m)$ can be found e.g. in [3] or [7]; $(l-1)$ -dimensional volume in a rational $(l-1)$ -dimensional affine subspace of $L_{\mathcal{I}}$ has to be normalized in such a way that the volume of the unit cube spanned by any integer basis of the corresponding linear subspace is equal to 1. Let us recall that $V_m(\underbrace{\Delta, \dots, \Delta}_{m \text{ terms}})$ is simply the m -dimensional volume of Δ . We have to assume that $V_0(\text{nothing}) = 1$, (this is necessary to define V_a for $l = 1$). Let:

$$\begin{aligned} \zeta_{\mathcal{I}}^0(t) &= \prod_{a \in E_{\mathcal{I}} : m(a, \Gamma_1) > m(a, \Gamma_2)} (1 - t^{m(a, \Gamma_1) - m(a, \Gamma_2)})^{(l-1)! V_a}, \\ \zeta_{\mathcal{I}}^\infty(t) &= \prod_{a \in E_{\mathcal{I}} : m(a, \Gamma_1) < m(a, \Gamma_2)} (1 - t^{m(a, \Gamma_2) - m(a, \Gamma_1)})^{(l-1)! V_a}, \\ \zeta_l^\bullet(t) &= \prod_{\mathcal{I} : \#\mathcal{I} = l} \zeta_{\mathcal{I}}^\bullet(t), \\ \zeta_\Lambda^\bullet(t) &= \prod_{l=1}^{n+1} (\zeta_l^\bullet(t))^{(-1)^{l-1}}, \end{aligned}$$

where $\bullet = 0$ or ∞ .

THEOREM 3 *Let $f = \frac{P}{Q}$ be a germ of a meromorphic function on $(\mathbb{C}^{n+1}, 0)$ non-degenerated with respect to its Newton pair $\Lambda = (\Gamma_1, \Gamma_2)$. Then*

$$\zeta_f^0(t) = \zeta_\Lambda^0(t) \quad \text{and} \quad \zeta_f^\infty(t) = \zeta_\Lambda^\infty(t).$$

Proof. Let Σ be an unimodular simplicial subdivision of $\mathbb{R}_{\geq 0}^{n+1}$ which corresponds to the pair (Γ_1, Γ_2) of Newton diagrams in the sense of [7] Section 4. This subdivision is consistent with each of the Newton diagrams Γ_1 and Γ_2 in the sense of [8].

Let $\pi : (\mathcal{X}, \mathcal{D}) \rightarrow (\mathbb{C}^{n+1}, 0)$ be the toroidal modification map corresponding to Σ . Since the pair (P, Q) is non-degenerated with respect to (Γ_1, Γ_2) , π is a resolution of the germ $f = \frac{P}{Q}$ (see [7]). We have the sets $S_{k,l} = S_k(P) \cap S_l(Q)$. The description of $S_k(P)$ (and of $S_l(Q)$) can be found in [8], Section 7. Each of them consists of open parts of certain complex tori of some dimensions.

Tori of dimension n correspond to one-dimensional cone of Σ which are positive (i.e., lie in $(\mathbb{R}^*)_+^{n+1}$). The multiplicity of $P \circ \pi$ (respectively of $Q \circ \pi$) at such a torus is equal to $m(a, \Gamma_1)$, (respectively to $m(a, \Gamma_2)$) for the primitive integer covector a which spans the corresponding cone.

Tori of dimension $(l-1)$ correspond to positive simplicial $(n+2-l)$ -dimensional cones of Σ which have a cone of the form

$$\mathfrak{S} = \{a \in (\mathbb{R}^*)_{\geq 0}^{n+1} : a_j > 0 \text{ for } j \notin \mathcal{I}, \quad a_j = 0 \text{ for } j \in \mathcal{I}\}$$

with $\#(\mathcal{I}) = l$ (these cones are elements of Σ) as its face. Moreover these cones correspond to one-dimensional cones of a partitioning of $L_{\mathcal{I}}$ which is consistent with the Newton diagrams $\Gamma_{i,\mathcal{I}} = \Gamma_i \cap L_{\mathcal{I}} \subset L_{\mathcal{I}}$. The multiplicities of $P \circ \pi$ and $Q \circ \pi$ at such a torus again are equal to $m_{\mathcal{I}}(a, \Gamma_{1,\mathcal{I}})$ and $m_{\mathcal{I}}(a, \Gamma_{2,\mathcal{I}})$ for the primitive integer covector a from the corresponding one-dimensional cone.

In order to apply Theorem 1 we have to calculate the Euler characteristic of the corresponding part of an $l-1$ -dimensional torus T : the complement to the intersection with the strict transform of the hypersurface $H = \{P = 0\} \cup \{Q = 0\}$. Let A (respectively B) the intersection of the torus T with the strict transform of the hypersurface $\{P = 0\}$ (respectively of $\{Q = 0\}$), let $\Delta_i := \Delta(a, \Gamma_{i,\mathcal{I}})$. From the results of Khovanskii ([6]) it follows that the Euler characteristic of A (respectively of B) is equal to $(-1)^l(l-1)!V_{l-1}(\underbrace{\Delta_1, \dots, \Delta_1}_{l-1 \text{ terms}})$

(respectively to $(-1)^l(l-1)!V_{l-1}(\underbrace{\Delta_2, \dots, \Delta_2}_{l-1 \text{ terms}})$), the Euler characteristic of $A \cap B$ is equal to

$$(-1)^{l-1}(l-1)![V_{l-1}(\underbrace{\Delta_1, \dots, \Delta_1}_{l-2 \text{ terms}}, \Delta_2) + V_{l-1}(\underbrace{\Delta_1, \dots, \Delta_1}_{l-3 \text{ terms}}, \Delta_2, \Delta_2) + \dots + V_{l-1}(\Delta_1, \underbrace{\Delta_2, \dots, \Delta_2}_{l-2 \text{ terms}})].$$

Thus the Euler characteristic of the complement of $A \cup B$ in the torus T is equal to

$$\begin{aligned} & \chi(T) - \chi(A) - \chi(B) + \chi(A \cap B) = \\ & = (-1)^{l-1} (l-1)! [V_{l-1}(\underbrace{\Delta_1, \dots, \Delta_1}_{l-1 \text{ terms}}) + V_{l-1}(\underbrace{\Delta_1, \dots, \Delta_1}_{l-2 \text{ terms}}, \Delta_2) + \dots + V_{l-1}(\underbrace{\Delta_2, \dots, \Delta_2}_{l-1 \text{ terms}})], \end{aligned}$$

which implies the statement. $\square$

5 The Varchenko type formula for $f = \frac{P}{z_0^d}$

As we have mentioned at the beginning, in order to study the behaviour of polynomials at infinity, germs of meromorphic functions of the form $\frac{P(z_0, z_1, \dots, z_n)}{z_0^d}$ have to be of interest. In this case the formulae for the zeta-functions $\zeta_\Lambda^0(t)$ and $\zeta_\Lambda^\infty(t)$ are considerably reduced. Thus let us reformulate the definition of these zeta-functions for the case when the Newton diagram Γ_2 consists of one point $(d, 0, \dots, 0)$ (in terms of the Newton diagram $\Gamma := \Gamma_1$ of P). The description is as follows.

Let $1 \leq l \leq n+1$ and let $\mathcal{I}$ be a subset of $\{1, \dots, n\}$ with the number of elements $\#\mathcal{I}$ equal to $l-1$. Let $\gamma_1^\mathcal{I}, \dots, \gamma_{j(\mathcal{I})}^\mathcal{I}$ be all $(l-1)$ -dimensional faces of $\Gamma_{\mathcal{I} \cup \{0\}}$ and $a_{\mathcal{I},1}, \dots, a_{\mathcal{I},j(\mathcal{I})}$ the corresponding primitive covectors (normal to $\gamma_1^\mathcal{I}, \dots, \gamma_{j(\mathcal{I})}^\mathcal{I}$), $a_{\mathcal{I},s}^0$ is the 0th coordinate of $a_{\mathcal{I},s}$, let $m_s(\mathcal{I}) = (a_{\mathcal{I},s}, k)$ for $k \in \gamma_s^\mathcal{I}$. Then

$$\begin{aligned} \zeta_{\mathcal{I} \cup \{0\}}^0(t) &= \prod_{1 \leq s \leq j(\mathcal{I}): m_s(\mathcal{I}) > d \cdot a_{\mathcal{I},s}^0} (1 - t^{m_s(\mathcal{I}) - d \cdot a_{\mathcal{I},s}^0})^{(l-1)! V_{l-1}(\gamma_s^\mathcal{I})}, \\ \zeta_{\mathcal{I} \cup \{0\}}^\infty(t) &= \prod_{1 \leq s \leq j(\mathcal{I}): m_s(\mathcal{I}) < d \cdot a_{\mathcal{I},s}^0} (1 - t^{d \cdot a_{\mathcal{I},s}^0 - m_s(\mathcal{I})})^{(l-1)! V_{l-1}(\gamma_s^\mathcal{I})}, \\ \zeta_l^\bullet(t) &= \prod_{\mathcal{I} \subset \{1, \dots, n\}: \#\mathcal{I} = l-1} \zeta_{\mathcal{I} \cup \{0\}}^\bullet(t), \\ \zeta_\Lambda^\bullet(t) &= \prod_{l=1}^{n+1} (\zeta_l^\bullet(t))^{(-1)^{l-1}}. \end{aligned}$$

($\bullet = 0$ or ∞) where $V_{l-1}(\gamma_s^\mathcal{I})$ is the (usual) $(l-1)$ -dimensional volume of the face $\gamma_s^\mathcal{I}$ (in the hyperplane spanned by it in $L_{\mathcal{I} \cup \{0\}}$).

6 Examples

Example 1. Let $f = \frac{x^3 - xy}{y}$. The Milnor fibre $\mathcal{M}_f^0$ (repectively $\mathcal{M}_f^\infty$) is $\{(x, y) : \|(x, y)\| < \varepsilon, (x^3 - xy, y) \in B_\delta, x^3 - xy = cy\} \setminus \{(0, 0)\}$, where $\|c\|$ is small (repectively large). From the equation $x^3 - xy = cy$ one has $y = \frac{x^3}{x+c}$ and thus $\mathcal{M}_f^0$ is diffeomorphic to the disk $\mathcal{D}$ in the x -plane with two points removed: $-c$ and the origin. In the same way $\mathcal{M}_f^\infty$ is

diffeomorphic to the punctured disk $\mathcal{D}^*$. It is not difficult to understand that the action of the monodromy transformation in the homology groups is trivial in both cases. Thus

$$\zeta_f^0(t) = (1-t)^{-1} \quad \text{and} \quad \zeta_f^\infty(t) = 1.$$

Now let us calculate these zeta functions from their Newton diagrams, Fig 1.




Figure 1.

We have $\zeta_1^\bullet(t) = 1$ since each coordinate axis intersects only one Newton diagram. There is only one linear function (namely $a = k_x + 2k_y$) such that $\dim \Delta(a, \Gamma_1) = 1$. The one-dimensional volume $V_1(\Delta(a, \Gamma_1))$ of $\Delta(a, \Gamma_1)$ is equal to 1 and $V_1(\Delta(a, \Gamma_2)) = 0$. We have $m(a, \Gamma_1) = 3$ and $m(a, \Gamma_2) = 2$. Thus $\zeta_2^0(t) = (1-t)$, $\zeta_2^\infty(t) = 1$, $\zeta_{(\Gamma_1, \Gamma_2)}^0(t) = (1-t)^{-1}$ and $\zeta_{(\Gamma_1, \Gamma_2)}^\infty(t) = 1$ which coincides with the formulae for f written above.

Example 2. Let $P = xyz + x^p + y^q + z^r$ be a $T_{p,q,r}$ singularity, $\frac{1}{p} + \frac{1}{q} + \frac{1}{r} < 1$ and let $Q = x^d + y^d + z^d$ be a homogeneous polynomial of degree d . Suppose that $p > q > r > d > 3$ and that p, q , and r are pairwise prime. Let us compute the zeta-functions of $f = \frac{P}{Q}$ using Theorems 2 and 3.

(a) It is clear that f is non-degenerated with respect to its Newton pair $\Lambda = (\Gamma_1, \Gamma_2)$. Thus

$$\zeta_f^\bullet(t) = \zeta_\Lambda^\bullet(t) = \zeta_1^\bullet(\zeta_2^\bullet)^{-1} \zeta_3^\bullet \quad (\bullet = 0 \text{ or } \infty).$$

One has $\zeta_1^\infty = \zeta_2^\infty = 1$ and the unique covector which is necessary for computing ζ_3^∞ is $a = (1, 1, 1)$. In this case $m(a, \Gamma_1) = 3$, $m(a, \Gamma_2) = d$, $\Delta(a, \Gamma_1) = \{(1, 1, 1)\}$ and $\Delta(a, \Gamma_2)$ is the simplex $\{k_x + k_y + k_z = d, k_x \geq 0, k_y \geq 0, k_z \geq 0\}$, its two-dimensional volume is equal to $\frac{d^2}{2}$. Thus $\zeta_f^\infty = (1 - t^{d-3})^{d^2}$.

We have

$$\begin{aligned} \zeta_1^0 &= (1 - t^{p-d})(1 - t^{q-d})(1 - t^{r-d}), \\ \zeta_2^0 &= (1 - t^{r(q-d)})(1 - t^{r(p-d)})(1 - t^{q(p-d)})(1 - t^{r-d})^{2d}(1 - t^{q-d})^d. \end{aligned}$$

To compute ζ_3^0 one has to take into account both covectors $(rq - q - r, r, q)$, $(r, pr - p - r, p)$, and $(q, p, qp - p - q)$, corresponding to two-dimensional faces of Γ_1 , and covectors $(1, r - 2, 1)$, $(r - 2, 1, 1)$, and $(q - 2, 1, 1)$, corresponding to pairs of the form (one-dimensional face of Γ_1 , one-dimensional face of Γ_2). E.g., for $a = (1, r - 2, 1)$, $\Delta(a, \Gamma_1)$ (respectively $\Delta(a, \Gamma_2)$) is the segment between $(0, 0, r)$ to $(1, 1, 1)$ (respectively between

$(d, 0, 0)$ and $(0, 0, d)$. Pay attention to the “absence of the symmetry”: last three covectors are not obtained from each other by permutting the coordinates and the numbers p , q , and r . This way

$$\zeta_3^0 = (1 - t^{r(q-d)})(1 - t^{r(p-d)})(1 - t^{q(p-d)})(1 - t^{r-d})^{2d}(1 - t^{q-d})^d$$

and

$$\zeta_f^0 = (1 - t^{p-d})(1 - t^{q-d})(1 - t^{r-d}).$$

(b) For computing the zeta-functions of f with the help of Theorem 2, let $\pi : (\mathcal{X}, \mathcal{D}) \rightarrow (\mathbb{C}^3, 0)$ be the blowing-up of the origin in $\mathbb{C}^3$ and let φ be the lifting $f \circ \pi$ of f to the space $\mathcal{X}$. The exceptional divisor $\mathcal{D}$ is the complex projective plane $\mathbb{CP}^2$. Let H_1 and H_2 be the strict transforms of the hypersurfaces $\{P = 0\}$ and $\{Q = 0\}$, $D_i = \mathcal{D} \cap H_i$. The curve D_1 consists of three transversal lines l_1, l_2, l_3 and has three singular points $S_1 = l_2 \cap l_3 = (0, 0, 1)$, $S_2 = l_1 \cap l_3 = (0, 1, 0)$, and $S_3 = l_1 \cap l_2 = (1, 0, 0)$. The curve D_2 is a smooth curve of degree d , it intersects D_1 at $3d$ different points $\{P_1, \dots, P_{3d}\}$.

One has the following natural stratification of the exceptional divisor $\mathcal{D}$:

- (i) 0-dimensional strata Λ_i^0 ($i = 1, 2, 3$), each consisting of one point S_i ;
- (ii) 0-dimensional strata Ξ_i^0 consisting of one point P_i each ($i = 1, \dots, 3d$);
- (iii) 1-dimensional strata $\Xi_i^1 = l_i \setminus \{D_2 \cup l_j \cup l_k\}$ ($i = 1, 2, 3$) and $\Xi_4^1 = D_2 \setminus D_1$;
- (iv) 2-dimensional stratum $\Xi^2 = \mathcal{D} \setminus (D_1 \cup D_2)$.

It is not difficult to see that $\zeta_{\Xi_2^0}(t) = 1$, $\zeta_{\Xi_2^\infty}(t) = 1 - t^{d-3}$, for each stratum Ξ from Ξ_i^0 ($1 \leq i \leq 3d$), Ξ_i^1 ($1 \leq i \leq 4$) one has $\zeta_\bullet^\bullet(t) = 1$ ($\bullet = 0$ or ∞).

In what follows the exceptional divisor $\mathcal{D}$ has the local equation $u = 0$. At the point S_1 the lifting φ of the function f is of the form $\frac{u^3 x_1 y_1 + u^r + x_1^p u^p + y_1^q u^q}{u^d x_1^d + u^d y_1^d + u^d}$. This germ has the same Newton pair as the germ $\frac{u^3 x_1 y_1 + u^r}{u^d}$. Using theorem 3 one has $\zeta_{\Lambda_1^0}^\infty = 1$, $\zeta_{\Lambda_1^0}^\infty = 1 - t^{r-d}$. At the point S_2 the function φ has the form $\frac{u^3 x_1 z_1 + z_1^r u^r + x_1^p u^p + u^q}{u^d x_1^d + u^d + z_1^d u^d}$. It has the same Newton pair as $\frac{u^3 x_1 z_1 + z_1^r u^r + u^q}{u^d}$. Using Theorem 3 one has $\zeta_{\Lambda_2^0}^\infty(t) = 1$, $\zeta_{\Lambda_2^0}^0(t) = 1 - t^{q-d}$. Just in the same way $\zeta_{\Lambda_3^0}^\infty(t) = 1$, $\zeta_{\Lambda_3^0}^0(t) = 1 - t^{p-d}$. Combining these computations together, one has the same results as above (without using a partial resolution).

References

- [1] A'CAMPO N., La fonction zeta d'une monodromie, *Comment. Math. Helv.* 50 (1975), 233-248.
- [2] ARNOLD V.I., GUSEIN-ZADE S.M., VARCHENKO A.N., *Singularities of Differentiable Maps, Vol. II*, (Birkhäuser, Boston-Basel- Berlin. 1988).

- [3] BERNSHTEIN D.N., The number of roots of a system of equations, *Funct. Anal. Appl.* 9 (1975), no. 3, 1-4.
- [4] GUSEIN-ZADE S.M., LUENGO I., MELLE-HERNÁNDEZ A. Partial resolutions and the zeta-function of a singularity, to be published in *Comment. Math. Helv.*
- [5] KHOVANSKII A.G., Newton polyhedra and toroidal varieties, *Funct. Anal. Appl.* 11 (1977), no. 4, 56-64.
- [6] KHOVANSKII A.G., Newton polyhedra and the genus of complete intersections, *Funct. Anal. Appl.* 12 (1978), no. 1, 51-61.
- [7] OKA M., Principal zeta-function of non-degenerate complete intersection singularity, *J. Fac. Sci. Uni. Tokyo Sect. IA, Math.* 37 (1990), 11-32.
- [8] VARCHENKO A.N., Zeta function of monodromy and Newton's diagram, *Invent. Math.*, 37 (1976), 253-262.

Addresses:

Sabir M. Gusein-Zade,
 Moscow State University,
 Faculty of Mathematics and Mechanics.
 Moscow, 119899, Russia.
 e-mail: sabir@ium.ips.ras.ru

Ignacio Luengo,
 Departamento de Algebra,
 Facultad de Ciencias Matemáticas,
 Universidad Complutense de Madrid.
 Ciudad Universitaria, 28040 Madrid, Spain.
 e-mail: iluengo@eucmos.sim.ucm.es

Alejandro Melle-Hernández,
 Departamento de Geometría y Topología,
 Facultad de Ciencias Matemáticas,
 Universidad Complutense de Madrid.
 Ciudad Universitaria, 28040 Madrid, Spain.
 e-mail: amelle@eucmos.sim.ucm.es